

Non-Abelian Stokes Theorem and Quark Confinement in SU(3) Yang-Mills Gauge Theory*

Kei-Ichi Kondo^{1,2†} and Yutaro Taira^{2‡}

¹ *Department of Physics, Faculty of Science, Chiba University, Chiba 263-8522, Japan*

² *Graduate School of Science and Technology, Chiba University, Chiba 263-8522, Japan*

Abstract

We derive a new version of SU(3) non-Abelian Stokes theorem by making use of the coherent state representation on the coset space $SU(3)/(U(1) \times U(1)) = F_2$, the flag space. Then we outline a derivation of the area law of the Wilson loop in SU(3) Yang-Mills theory in the maximal Abelian gauge (The detailed exposition will be given in a forthcoming article). This derivation is performed by combining the non-Abelian Stokes theorem with the reformulation of the Yang-Mills theory as a perturbative deformation of a topological field theory recently proposed by one of the authors. Within this framework, we show that the fundamental quark is confined even if $G = SU(3)$ is broken by partial gauge fixing into $H = U(2)$ just as G is broken to $H = U(1) \times U(1)$. An origin of the area law is related to the geometric phase of the Wilczek-Zee holonomy for $U(2)$. Abelian dominance is an immediate byproduct of these results and magnetic monopole plays the dominant role in this derivation.

Key words: quark confinement, topological field theory, magnetic monopole, non-Abelian Stokes theorem, topological soliton

PACS: 12.38.Aw, 12.38.Lg

[†]E-mail: kondo@cuphd.nd.chiba-u.ac.jp

[‡]E-mail: taira@cuphd.nd.chiba-u.ac.jp

* To be published in Mod. Phys. Lett. A.

1 Introduction

Quantum Chromodynamics (QCD) is a quantized gauge field theory with color $SU(3)$ as the gauge group. Usually, a simplified version of QCD with a gauge group $SU(2)$ is first investigated to avoid technical complexity of dealing with $SU(3)$ group. This is also the case in the study of quark confinement in QCD. However, this simplification might lose some important features which may appear only when we begin to consider $SU(3)$ case. This is anticipated because it is generally believed that the quark confinement is not entirely of the kinematic origin and that some dynamical information on non-Abelian Yang-Mills gauge theory is indispensable to lead to the complete resolution of the problem.

Recently, the dual superconductor picture [1] of quark confinement in QCD has been extensively investigated based on the idea of Abelian projection [2], see [3] for review. In this scenario, the original gauge group $G = SU(N)$ is broken down to the maximal torus subgroup $H = U(1)^{N-1}$ (up to the discrete Weyl symmetry) by the partial gauge fixing. This will imply that the $(N - 1)$ species of Abelian magnetic monopoles will be responsible for quark confinement ($N \geq 2$). In the presence of the elementary scalar field, e.g., in the Yang-Mills-Higgs theory, we can break the gauge symmetry G to any subgroup H by choosing appropriately the expectation value of the Higgs scalar field. However, QCD does not have the elementary scalar field in the theory, in sharp contrast with the supersymmetric version [4]. If the scalar field would appear in QCD and carry the degrees of freedom of magnetic monopoles, it should be provided as a composite of gauge field. This was indeed the case in $SU(2)$, see [5, 6]. The dynamics of true $SU(3)$ QCD may restrict the type of the composite scalar field as a carrier of magnetic monopoles which is responsible for the quark confinement. If so, the subgroup H could be restricted.

In this Letter, we give a first report on the extension of the previous analyses of quark confinement in $SU(2)$ Yang-Mills theory performed in a series of papers [5, 6, 7, 8] to the $SU(3)$ case. The technical details will be given in a subsequent

paper [9]. Our results suggest that the fundamental quark in QCD is confined in the sense of the area law of the Wilson loop even if $SU(3)$ is restricted to $H = U(2)$ just as $U(1) \times U(1)$, without contradicting with the maximal Abelian gauge.

2 Coherent state on the flag space

2.1 General procedure

In this Letter we shall derive a version of the non-Abelian Stokes theorem (NAST) for $SU(3)$, generalizing the NAST for $SU(2)$ derived in [10, 7]. Further generalization to $SU(N)$ is straightforward [9]. First of all, we construct the coherent state corresponding to the coset representatives $\xi \in G/\tilde{H}$. For inputs, we prepare (a) the gauge group G and its (semi simple) Lie algebra $\mathcal{G}$ with the generators $\{T^A\}$ of the Lie algebra being rewritten in terms of the Cartan basis $\{H_i, E_\alpha, E_{-\alpha}\}$; (b) the Hilbert space V^Λ as a carrier (the representation space) of the unitary irreducible representation Γ^Λ of G ; (c) a reference state $|\Lambda\rangle$ within the Hilbert space V^Λ , which can be normalized to unity, $\langle\Lambda|\Lambda\rangle = 1$.

We define the *maximal stability subgroup (isotropy subgroup)* $\tilde{H}$ as a subgroup of G that consists of all the group elements h that leave the reference state $|\Lambda\rangle$ invariant up to a phase factor, i.e., $h|\Lambda\rangle = |\Lambda\rangle e^{i\phi(h)}$, $h \in \tilde{H}$. The phase factor is unimportant in the following because we consider the expectation value of any operators in the coherent state. For every element $g \in G$, there is a unique decomposition of g into a product of two group elements, $g = \xi h$, $\xi \in G/\tilde{H}$, $h \in \tilde{H}$, for $g \in G$. We can obtain a unique coset space for a given $|\Lambda\rangle$. The action of arbitrary group element $g \in G$ on $|\Lambda\rangle$ is given by $g|\Lambda\rangle = \xi h|\Lambda\rangle = \xi|\Lambda\rangle e^{i\phi(h)}$.

The coherent state is constructed as $|\xi, \Lambda\rangle = \xi|\Lambda\rangle$. This definition of the coherent state is in one-to-one correspondence with the coset space $G/\tilde{H}$ and the coherent states preserve all the algebraic and topological properties of the coset space $G/\tilde{H}$. If $\Gamma^\Lambda(\mathcal{G})$ is Hermitian, then $H_i^\dagger = H_i$, and $E_\alpha^\dagger = E_{-\alpha}$. Thus the coherent state is given

by [11]

$$|\xi, \Lambda\rangle = \exp \left[\sum_{\beta'} (\eta_{\beta} E_{\beta} - \eta_{\beta}^* E_{-\beta}) \right] |\Lambda\rangle, \quad (1)$$

where $|\Lambda\rangle$ is the highest-weight state such that $|\Lambda\rangle$ is (i) annihilated by all the shift-up operators E_{α} with $\alpha > 0$, i.e., $E_{\alpha}|\Lambda\rangle = 0(\alpha > 0)$; (ii) mapped into itself by all diagonal operators H_i , i.e., $H_i|\Lambda\rangle = \Lambda_i|\Lambda\rangle$; (iii) annihilated by some shift-down operators E_{α} with $\alpha < 0$, not by other E_{β} with $\beta < 0$, i.e., $E_{\alpha}|\Lambda\rangle = 0(\text{some } \alpha < 0)$; $E_{\beta}|\Lambda\rangle = |\Lambda + \beta\rangle(\text{some } \beta < 0)$; and the sum $\sum_{\beta'}$ is restricted to those shift operators E_{β} which obey (iii).

Two coherent states are non-orthogonal, $\langle \xi', \Lambda | \xi, \Lambda \rangle \neq 0$, but normalized to unity, $\langle \xi, \Lambda | \xi, \Lambda \rangle = 1$. The coherent state spans the entire space V^{Λ} . By making use of the group-invariant measure $d\mu(\xi)$ of G which is appropriately normalized, we obtain

$$\int |\xi, \Lambda\rangle d\mu(\xi) \langle \xi, \Lambda| = I, \quad (2)$$

which shows that the coherent states are complete, but overcomplete. This resolution of identity is very important to obtain the path integral formula given below.

For concreteness, we focus on the $SU(3)$ case. Using the Dynkin index $[m, n]$ (m, n : integers), the highest weight Λ can be written as $\vec{\Lambda} = m\vec{h}_1 + n\vec{h}_2$ (m, n are non-negative integers for the highest weight) where h_1, h_2 are highest weights of two fundamental representations of $SU(3)$ corresponding to $[1, 0], [0, 1]$ respectively, i.e., $\vec{h}_1 = (\frac{1}{2}, \frac{1}{2\sqrt{3}}), \vec{h}_2 = (\frac{1}{2}, \frac{-1}{2\sqrt{3}})$. Therefore, $\vec{\Lambda} = (\frac{m+n}{2}, \frac{m-n}{2\sqrt{3}})$. The maximal stability group $\tilde{H}$ is given by $\tilde{H} = U(2)$ if $m = 0$ or $n = 0$ (case (I)), whereas $\tilde{H}$ is the maximal torus group $H = U(1) \times U(1)$ if $m \neq 0$ and $n \neq 0$ (case (II)). Therefore, for the representation with the highest weight Λ , the coset $G/\tilde{H}$ is given by $SU(3)/U(2) = SU(3)/(SU(2) \times U(1)) = CP^2$ in the case (I) and $SU(3)/(U(1) \times U(1)) = F_2$ in the case (II). Here, CP^{N-1} is the complex projective space and F_{N-1} is the flag space [12]. Therefore, the two fundamental representations belong to the case (I), so the maximal stability group is $\tilde{H} = U(2)$, rather than the maximal torus group $H = U(1) \times U(1)$.

The implications of this fact to the mechanism of quark confinement is discussed in the following.

2.2 Explicit form of the coherent state on flag space

The $SU(2)$ case is well known. The coherent state for $F_1 := SU(2)/U(1)$ is obtained as

$$|j, w\rangle = \xi(w)|j, -j\rangle = e^{\zeta J_+ - \bar{\zeta} J_-}|j, -j\rangle = \frac{1}{(1 + |w|^2)^j} e^{w J_+} |j, -j\rangle, \quad (3)$$

where $J_+ = J_1 + iJ_2$, $J_- = J_1^\dagger$, $|j, -j\rangle$ is the lowest state, $|j, m = -j\rangle$ of $|j, m\rangle$, and $w = \frac{\zeta \sin|\zeta|}{|\zeta| \cos|\zeta|}$. Note that $(1 + |w|^2)^{-j}$ is a normalization factor to ensure $\langle j, w | j, w \rangle = 1$, which is obtained from the Baker-Campbell-Hausdorff (BCH) formulas. The invariant measure is given by

$$d\mu = \frac{2j+1}{4\pi} \frac{dw d\bar{w}}{(1 + |w|^2)^2}. \quad (4)$$

For $J_A = \frac{1}{2}\sigma^A$ ($A = 1, 2, 3$) with Pauli matrices σ^A , we obtain $J_+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, and $e^{w J_+} = \begin{pmatrix} 1 & w \\ 0 & 1 \end{pmatrix} \in F_1 = CP^1 = SU(2)/U(1) \cong S^2$. The complex variable w is a CP^1 variable written as $w = e^{-i\phi} \tan \frac{\theta}{2}$, in terms of the polar coordinate on S^2 or Euler angles, see [6].

Now we proceed to the $SU(3)$ case. The coherent state for $F_2 = SU(3)/U(1)^2$ is given by $|\xi, \Lambda\rangle = \xi(w)|\Lambda\rangle := V^\dagger(w)|\Lambda\rangle$, with the highest(lowest)-weight state $|\Lambda\rangle$ and

$$|\xi, \Lambda\rangle = \exp \left[\sum_{\alpha=1}^3 (\zeta_\alpha E_\alpha - \bar{\zeta}_\alpha E_\alpha^\dagger) \right] |\Lambda\rangle = e^{-\frac{1}{2}K(w, \bar{w})} \exp \left[\sum_{\alpha=1}^3 \tau_\alpha E_\alpha \right] |\Lambda\rangle, \quad (5)$$

where $e^{-\frac{1}{2}K}$ is the normalization factor obtained from the Kähler potential (explained later)

$$K(w, \bar{w}) := \ln[(\Delta_1(w, \bar{w}))^m (\Delta_2(w, \bar{w}))^n], \quad (6)$$

$$\Delta_1(w, \bar{w}) := 1 + |w_1|^2 + |w_2|^2, \quad \Delta_2(w, \bar{w}) := 1 + |w_3|^2 + |w_2 - w_1 w_3|^2. \quad (7)$$

The coherent state $|\xi, \Lambda\rangle$ is normalized, so that $\langle \xi, \Lambda | \xi, \Lambda \rangle = 1$. We can obtain the inner product [9]

$$\langle \xi', \Lambda | \xi, \Lambda \rangle = e^{K(w, \bar{w}')} e^{-\frac{1}{2}[K(w', \bar{w}') + K(w, \bar{w})]}, \quad (8)$$

where $K(w, \bar{w}') := \ln[1 + \bar{w}_1' w_1 + \bar{w}_2' w_2]^m [1 + \bar{w}_3' w_3 + (\bar{w}_2' - \bar{w}_1' \bar{w}_3')(w_2 - w_1 w_3)]^n$.

The $SU(3)$ invariant measure is given (up to a constant factor) by

$$d\mu(\xi) = d\mu(w, \bar{w}) = D(m, n) [(\Delta_1)^m (\Delta_2)^n]^{-2} \prod_{\alpha=1}^3 dw_\alpha d\bar{w}_\alpha, \quad (9)$$

where $D(m, n) = \frac{1}{2}(m+1)(n+1)(m+n+2)$ is the dimension of the representation. For

$E_1 := (\lambda_1 - i\lambda_2)/(2\sqrt{2})$, $E_2 := (\lambda_4 - i\lambda_5)/(2\sqrt{2})$, $E_3 := (\lambda_6 - i\lambda_7)/(2\sqrt{2})$, with the Gell-

Mann matrices $\lambda_A (A = 1, \dots, 8)$, we obtain $\exp \left[\sum_{\alpha=1}^3 \sqrt{2} \tau_\alpha E_\alpha \right] = \begin{pmatrix} 1 & w_1 & w_2 \\ 0 & 1 & w_3 \\ 0 & 0 & 1 \end{pmatrix}^t \in$

$F_2 = SU(3)/U(1)^2$. Two sets of three complex variables are related as $w_1 = \tau_1$, $w_2 =$

$\tau_2 + \tau_1 \tau_3 / 2$, $w_3 = \tau_3$, or conversely $\tau_1 = w_1$, $\tau_2 = w_2 - w_1 w_3 / 2$, $\tau_3 = w_3$.

Any element of $F_{N-1} = SU(N)/U(1)^{N-1}$ is written as an upper triangular matrix with $N(N-1)/2$ complex numbers. It is not difficult to extend the above results to $SU(N)$, see [9].

3 Non-Abelian Stokes theorem for $SU(3)$

For the infinitesimal deviation $\xi' = \xi + d\xi$ (which is sufficient to derive the path integral formula), we find from (5) and (8)

$$\langle \xi', \Lambda | \xi, \Lambda \rangle = \exp(i\omega + O((dw)^2)), \quad (10)$$

$$\omega(x) := \langle \Lambda | iV(x) dV^\dagger(x) | \Lambda \rangle = \langle \Lambda | i\xi^\dagger(x) d\xi(x) | \Lambda \rangle, \quad (11)$$

where $d := dx^\mu \partial_\mu$ denotes an exterior derivative and the one-form ω is given by

$$\omega = im \frac{w_1 d\bar{w}_1 + w_2 d\bar{w}_2}{\Delta_1(w, \bar{w})} + in \frac{w_3 d\bar{w}_3 + (w_2 - w_1 w_3)(d\bar{w}_2 - \bar{w}_1 d\bar{w}_3 - \bar{w}_3 d\bar{w}_1)}{\Delta_2(w, \bar{w})}, \quad (12)$$

up to the total derivative. Here the x -dependence of V comes through that of $w(x)$ (the local field variable $w(x)$), i.e., $V(x) = V(w(x))$.

The Wilson loop operator $W^C[\mathcal{A}]$ is defined as the path-ordered exponent along the closed loop C , $W^C[\mathcal{A}] := \text{tr}[\mathcal{P} \exp(ig \oint_C \mathcal{A})] / \text{tr}(1)$, where $\mathcal{A}$ is the Lie-algebra valued connection one-form, $\mathcal{A}(x) = \mathcal{A}_\mu^A(x) T^A dx^\mu = \mathcal{A}^A(x) T^A$. Repeating the same

steps as those given in [7], we obtain the path integral representation of the Wilson loop,

$$\begin{aligned} W^C[\mathcal{A}] &= \int [d\mu(\xi)]_C \exp \left(ig \oint_C \langle \Lambda | [V \mathcal{A} V^\dagger + \frac{i}{g} V dV^\dagger] | \Lambda \rangle \right) \\ &= \int [d\mu(\xi)]_C \exp \left(ig \oint_C [n^A \mathcal{A}^A + \frac{1}{g} \omega] \right), \end{aligned} \quad (13)$$

where $[d\mu(\xi)]_C$ is the product measure of $d\mu(w(x), \bar{w}(x))$ along the loop, and

$$n^A(x) := \langle \Lambda | V(x) T^A V^\dagger(x) | \Lambda \rangle. \quad (14)$$

Using the (usual) Stokes theorem $\oint_{C=\partial S} \omega = \int_S d\omega$, we arrive at the NAST for $SU(3)$:

$$W^C[\mathcal{A}] = \int [d\mu(\xi)]_C \exp \left(ig \int_{S:\partial S=C} [d(n^A \mathcal{A}^A) + \frac{1}{g} \Omega_K] \right), \quad \Omega_K := d\omega. \quad (15)$$

Taking into account $d = \partial + \bar{\partial} = dw_\alpha \frac{\partial}{\partial w_\alpha} + d\bar{w}_\beta \frac{\partial}{\partial \bar{w}_\beta}$, we find for $SU(3)$

$$\begin{aligned} \Omega_K &= d\omega = im(\Delta_1)^{-2}[(1 + |w_1|^2)dw_2 \wedge d\bar{w}_2 - \bar{w}_2 w_1 dw_2 \wedge d\bar{w}_1 \\ &\quad - w_2 \bar{w}_1 dw_1 \wedge d\bar{w}_2 + (1 + |w_2|^2)dw_1 \wedge d\bar{w}_1] \\ &\quad + in(\Delta_2)^{-2}[\Delta_1 dw_3 \wedge d\bar{w}_3 - (w_1 + \bar{w}_3 w_2)dw_3 \wedge (d\bar{w}_2 - \bar{w}_3 d\bar{w}_1) \\ &\quad - (\bar{w}_1 + w_3 \bar{w}_2)(dw_2 - w_3 dw_1) \wedge d\bar{w}_3 \\ &\quad + (1 + |w_3|^2)(dw_2 - w_3 dw_1)(d\bar{w}_2 - \bar{w}_3 d\bar{w}_1)]. \end{aligned} \quad (16)$$

For $SU(2)$, we reproduce the well-known results; $\omega = im \frac{w d\bar{w}}{1+|w|^2}$ and $\Omega_K = im(1 + |w|^2)^{-2} dw \wedge d\bar{w}$.

The flag manifold F_{N-1} (including CP^{N-1}) is a Kähler manifold. So, it possesses complex *local* coordinates w_α , an Hermitian Riemannian metric $ds^2 = g_{\alpha\bar{\beta}} dw^\alpha d\bar{w}^\beta$, and a corresponding two-form, $\Omega_K = ig_{\alpha\bar{\beta}} dw^\alpha \wedge d\bar{w}^\beta$ (Kähler form) which is closed, i.e., $d\Omega_K = 0$. Any closed form Ω_K is *locally* exact, $\Omega_K = d\omega$ due to Poincaré's lemma, which is consistent with (16). The metric $g_{\alpha\bar{\beta}}$ is obtained as $g_{\alpha\bar{\beta}} = \frac{\partial}{\partial w^\alpha} \frac{\partial}{\partial \bar{w}^\beta} K$, from the Kähler potential $K = K(w, \bar{w})$. Indeed, the Ω_K just obtained in (16) agrees with the Kähler two-form, $\Omega_K = i\partial\bar{\partial}K$ obtained from the Kähler potential for F_2 , $K(w, \bar{w}) = \ln[(\Delta_1)^m (\Delta_2)^n]$. Hence, $\omega = \frac{i}{2}(\partial - \bar{\partial})K$, since $\partial^2 = 0 = \bar{\partial}^2, \partial\bar{\partial} + \bar{\partial}\partial = 0$.

For CP^2 , the Kähler potential $K(w, \bar{w}) = \ln[(\Delta_1)^m]$ is obtained as a special case of F_2 by putting $w_3 = 0$ and $n = 0$. For $F_1 = CP^1$, $K(w, \bar{w}) = \ln[(1 + |w|^2)^m]$, $m = 2j$.

We define $n^A(x) := \langle \Lambda | V(x) T^A V^\dagger(x) | \Lambda \rangle = \frac{1}{2} \bar{\omega}_a(x) (\lambda^A)_{ab} \omega_b(x)$, where $\omega_a(x) := V^\dagger(x) | \Lambda \rangle$. Especially, in the CP^2 case, we can write

$$n^A(x) := \langle \Lambda | U(x) T^A U^\dagger(x) | \Lambda \rangle = \frac{1}{2} \phi_a^*(x) (\lambda^A)_{ab} \phi_b(x), \quad (17)$$

where $U \in SU(3)$ and $|\Lambda\rangle = (1, 0, 0)^t$. Then $\phi_a := (U^\dagger(x) | \Lambda \rangle)_a = U_{1a}^*$ and hence $n^A = \frac{1}{2} (U \lambda^A U^\dagger)_{11}$. On the other hand, we examine another expression (adjoint orbit representation),

$$n^A = \text{tr}(U^\dagger \mathcal{H} U T^A) = \frac{1}{2} \text{tr}(\mathcal{H} U \lambda^A U^\dagger), \quad \mathbf{n} := n^A T^A = U^\dagger \mathcal{H} U, \quad (18)$$

where we take $\mathcal{H} = \vec{\Lambda} \cdot (\lambda^3, \lambda^8) = \Lambda_1 \lambda^3 + \Lambda_2 \lambda^8$. For $[1, 0]$ or $[0, 1]$, we find $\mathcal{H} = \text{diag}(\frac{2}{3}, -\frac{1}{3}, -\frac{1}{3})$, or $\text{diag}(\frac{1}{3}, -\frac{2}{3}, \frac{1}{3})$, and two definitions (17) and (18) are equivalent, $n^A = \frac{1}{2} \text{tr}(\mathcal{H} U \lambda^A U^\dagger) = \frac{1}{2} (U \lambda^A U^\dagger)_{11}$, since $U \lambda^A U^\dagger$ is traceless. The F_2 variables w_a and the CP^2 variables ϕ_a are related as $\phi_1 = 1, \phi_2 = w_1, \phi_3 = w_2$ and $\omega_a = \phi_a / \phi_1 = w_{a-1}$ ($w_0 := 1$ by definition). In the CP^{N-1} case, $\langle \Lambda | f(V) | \Lambda \rangle = 2 \text{tr}[\mathcal{H} f(U)]$. Hence, $\omega(x) := 2 \text{tr}[\mathcal{H} i U(x) dU^\dagger(x)] = -i 2 \text{tr}[\mathcal{H} dU(x) U^\dagger(x)]$ which is a diagonal piece of the Maurer-Cartan one-form $dU U^{-1}$. The two-form Ω_K is the symplectic two-form [13], $\Omega_K = d\omega = 2 \text{tr}(\mathcal{H} [U^{-1} dU, U^{-1} dU]) = 2 \text{tr}(\mathbf{n} [d\mathbf{n}, d\mathbf{n}])$.

For $SU(3)$, the topological part $\int_S \Omega_K = \oint_C \omega$ corresponding to the residual $U(2)$ invariance is interpreted as the geometric phase of the Wilczek-Zee holonomy [14], just as it is interpreted in the $SU(2)$ case as the Berry-Aharonov-Anandan phase for the residual $U(1)$ invariance. The details will be given in a subsequent paper [9].

4 Abelian dominance and area law

The NAST (15) implies that the expectation value of the Wilson loop in the $SU(N)$ Yang-Mills theory is given by

$$\langle W^C[\mathcal{A}] \rangle_{YM} = \left\langle \exp \left(ig \int_{S: \partial S = C} [da] \right) \right\rangle_{YM} = \left\langle \exp \left(ig \int_S [d(n^A \mathcal{A}^A) + \frac{1}{g} \Omega_K] \right) \right\rangle_{YM}, \quad (19)$$

where the one-form a is written as

$$a := n^A \mathcal{A}^A + \frac{1}{g} \omega = \langle \Lambda | \mathcal{A}^V | \Lambda \rangle. \quad (20)$$

Here $\mathcal{A}^V := V \mathcal{A} V^\dagger + \frac{i}{g} V dV^\dagger$ is the gauge transformation of $\mathcal{A}$ by $V \in F_{N-1}$. For quark in the fundamental representation, $a = 2\text{tr}(\mathcal{H} \mathcal{A}^V)$. So, the one-form a is equal to the diagonal piece of the gauge-transformed potential $\mathcal{A}^V$. In the $SU(2)$ case, $a = \text{tr}(T^3 \mathcal{A}^V)$ for any representation and the two-form $f := da$ is the Abelian field strength which is invariant under the $SU(2)$ transformation. The two-form f is nothing but the 't Hooft tensor $f_{\mu\nu}$ describing the magnetic flux emanating from the magnetic monopole, $f_{\mu\nu}(x) := \partial_\mu(n^A(x) \mathcal{A}_\nu^A(x)) - \partial_\nu(n^A(x) \mathcal{A}_\mu^A(x)) - \frac{1}{g} \mathbf{n}(x) \cdot (\partial_\mu \mathbf{n}(x) \times \partial_\nu \mathbf{n}(x))$, if we identify n^A with the direction of the Higgs field, $\hat{\phi}^A := \phi^A/|\phi|$, $|\phi| := \sqrt{\phi^A \phi^A}$. In general, the (curvature) two-form $f = d(n^A \mathcal{A}^A) + \frac{1}{g} \Omega_K$ in the NAST is the Abelian field strength which is invariant even under the non-Abelian gauge transformation of G . It reduces to the 't Hooft tensor in $SU(2)$ case.

The Abelian dominance in $SU(N)$ Yang-Mills theory is derived as follows. We adopt the maximal Abelian (MA) gauge. The MA gauge for $SU(N)$ is defined as follows. Under the Cartan decomposition of $\mathcal{A}$ into the diagonal (H) and off-diagonal (G/H) pieces, $\mathcal{A} = a^i H^i + A^a T^a$, the MA gauge condition is given by $\partial_\mu A_\mu^a - g f^{abi} a_\mu^i A_\mu^b = 0$, which is obtained by minimizing the functional $\mathcal{R} := \int d^4x \frac{1}{2} A_\mu^a A_\mu^a := \int d^4x \frac{1}{2} \text{tr}_{G/H}(\mathcal{A}_\mu \mathcal{A}_\mu)$, under the gauge transformation. For $SU(3)$, $H^1 = \frac{\lambda^3}{2}$, $H^2 = \frac{\lambda^8}{2}$, $T^a = \frac{\lambda^a}{2}$ ($a = 1, 2, 4, 5, 6, 7$). The low-energy effective gauge theory of QCD has been derived in the MA gauge by integrating out the off-diagonal gauge fields (together with the ghost and anti-ghost fields) [15, 5]. Then the $SU(N)$ Yang-Mills theory has been reduced to the Abelian gauge theory with the gauge coupling g which runs according to the same renormalization-group beta function as the original $SU(N)$ Yang-Mills theory,

$$\langle \cdots \rangle_{YM} = Z_{YM}^{-1} \int \mathcal{D}a \exp(i S_{APEGT}[a]) \cdots, \quad (21)$$

$$S_{APEGT}[a] = - \int d^4x \frac{1}{4g^2(\mu)} (da, da), \quad (22)$$

$$\frac{1}{g(\mu)^2} = \frac{1}{g(\mu_0)^2} + \frac{b_0}{8\pi^2} \ln \frac{\mu}{\mu_0}, \quad b_0 = \frac{11n}{3} > 0, \quad (23)$$

up to the one-loop level.¹ This result [5] combined with the $SU(2)$ NAST [10, 7] implies the Abelian dominance in the low-energy region of $SU(2)$ QCD, as shown in [7]. By virtue of the NAST for $SU(3)$ just derived, Abelian dominance in $SU(3)$ Yang-Mills theory follows immediately from the same argument as above, if we identify the connection one-form a with the diagonal piece of the gauge potential $\mathcal{A}^V$ given by (20). The monopole dominance is more subtle. It was derived for $SU(2)$ in [7] by showing that the dominant contribution to the area law comes from the monopole piece alone, $\Omega_K = d\omega = 2\text{tr}(\mathbf{n}[d\mathbf{n}, d\mathbf{n}])$.

In the following derivation of quark confinement, the magnetic monopoles (equivalently, the instantons in the coset F_{N-1} NLSM) is expected to give the dominant configuration to the area law of the Wilson loop or the string tension. The maximal stability group $\tilde{H}$ should be distinguished from the residual gauge group H retained after the partial gauge fixing $G \rightarrow H$ which realizes the magnetic monopole. The existence of magnetic monopole is suggested from the non-trivial Homotopy groups $\pi_2(G/\tilde{H})$. In the case (II), $\pi_2(SU(3)/(U(1) \times U(1))) = \pi_1(U(1) \times U(1)) = \mathbf{Z} + \mathbf{Z}$. On the other hand, in the case (I) $\pi_2(SU(3)/U(2)) = \pi_1(U(2)) = \pi_1(SU(2) \times U(1)) = \pi_1(U(1)) = \mathbf{Z}$. Note that CP^{N-1} NLSM has only the local $U(1)$ invariance for any $N \geq 2$. It is this $U(1)$ invariance that corresponds to a kind of Abelian magnetic monopole appearing in the case (I). This situation should be compared with the $SU(2)$ case where the maximal stability group $\tilde{H}$ is always given by the maximal torus $H = U(1)$ irrespective of the representation. Therefore, the coset is given by $G/\tilde{H} = SU(2)/U(1) = F_1 = CP^1 \cong S^2$ for any representation and $\pi_2(SU(2)/U(1)) = \mathbf{Z}$. For $SU(3)$, our results suggest that the fundamental quarks are to be confined even when the residual gauge group H is given by $\tilde{H} = U(2)$ and $\pi_2(G/H) = \mathbf{Z}$, while the adjoint quark is related to the maximal torus $H = U(1) \times U(1)$ and $\pi_2(G/H) = \mathbf{Z} + \mathbf{Z}$.

¹ The result of [15, 5] for $SU(2)$ can be generalized to $SU(N)$ in the straightforward way, at least in one-loop level [16]. In the two-loop level, it is not trivial. The two-loop result will be given elsewhere [16].

This observation is in sharp contrast with the conventional treatment in which the $(N - 1)$ species of magnetic monopoles corresponding to the residual maximal torus group $U(1)^{N-1}$ of $G = SU(N)$ are taken into account on equal footing.

We adopt an special choice of the MA gauge for fixing the gauge also in the $SU(3)$ case in which the gauge fixing part is given by $S_{GF} = \int d^4x i\delta_B \bar{\delta}_B \text{tr}_{G/H} \left[\frac{1}{2} \Omega_\mu \Omega_\mu + iC\bar{C} \right]$ where $\Omega_\mu := \frac{i}{g} U \partial_\mu U^\dagger$, with the BRST δ_B and anti-BRST $\bar{\delta}_B$ transformations, see [6] for details. We reformulate the Yang-Mills theory as a perturbative deformation of the topological quantum field theory (TQFT) which is obtained from the gauge fixing part for the compact gauge variable [6, 8]. Repeating the similar arguments given in [6, 7, 8], we can show [9] that the area decay of the Wilson loop $\langle W^C[\mathcal{A}] \rangle_{YM}$ in the four-dimensional Yang-Mills theory is dominated by the diagonal Wilson loop $\langle \exp(i \int_S \Omega_K) \rangle_{TQFT_4}$ in the four-dimensional TQFT with an action S_{GF} which describes the magnetic monopole in four dimensions. When the Wilson loop is planar, it is shown that this expectation value is equal to the the instanton distribution $\langle \exp(i \int_S \Omega_K) \rangle_{NLSM_2}$ in the two-dimensional coset G/H NLSM with an action S_{NLSM} . This is due to dimensional reduction. The dimensional reduction leads to the two-dimensional coset $F_{N-1} = SU(N)/U(1)^{N-1}$ NLSM,

$$S_{NLSM} = \frac{4\pi}{g^2(\mu)} \int d^2x g_{\alpha\bar{\beta}} \frac{\partial w^\alpha}{\partial x_a} \frac{\partial \bar{w}^\beta}{\partial x_a} = \frac{4\pi}{g^2(\mu)} \int dz d\bar{z} g_{\alpha\bar{\beta}} \left(\frac{\partial w^\alpha}{\partial z} \frac{\partial \bar{w}^\beta}{\partial \bar{z}} + \frac{\partial w^\alpha}{\partial \bar{z}} \frac{\partial \bar{w}^\beta}{\partial z} \right), (24)$$

where $g(\mu)$ is the running Yang-Mills coupling constant (23) induced from the perturbative deformation piece in four dimensions. Thus the derivation of the area law of the Wilson loop in the four-dimensional Yang-Mills theory is reduced to that of the diagonal Wilson loop in the two-dimensional coset NLSM.

The area law of the Wilson loop is shown as follows. The finite action configuration of the coset NLSM is provided with the instanton solution. For instanton configuration, $S_{NLSM} = \frac{4\pi^2}{g^2} |Q|$, with a topological charge Q . It is known [12] that the topological charge Q of the instanton in the F_{N-1} NLSM ($N \geq 3$) is given by the

integral of the Kähler 2-form,

$$Q = \frac{1}{\pi} \int \Omega_K = \int_{\mathbf{R}^2} \frac{d^2x}{\pi} \epsilon_{ab} g_{\alpha\bar{\beta}} \frac{\partial w^\alpha}{\partial x_a} \frac{\partial \bar{w}^\beta}{\partial x_b} = \int_C \frac{dz d\bar{z}}{\pi} g_{\alpha\bar{\beta}} \left(\frac{\partial w^\alpha}{\partial z} \frac{\partial \bar{w}^\beta}{\partial \bar{z}} - \frac{\partial w^\alpha}{\partial \bar{z}} \frac{\partial \bar{w}^\beta}{\partial z} \right). \quad (25)$$

This is a generalization of the F_1 case,

$$Q = \frac{i}{2\pi} \int_C \frac{dw d\bar{w}}{(1 + |w|^2)^2} = \frac{i}{2\pi} \int_{S^2} \frac{dz d\bar{z}}{(1 + |w|^2)^2} \left(\frac{\partial w}{\partial z} \frac{\partial \bar{w}}{\partial \bar{z}} - \frac{\partial w}{\partial \bar{z}} \frac{\partial \bar{w}}{\partial z} \right). \quad (26)$$

Since the Kähler potential for F_n is written as

$$K(w, \bar{w}) = \sum_{\ell=1}^n d_\ell K_\ell(w, \bar{w}) = \sum_{\ell=1}^n d_\ell \ln \Delta_\ell(w, \bar{w}) \quad (27)$$

with the Dynkin indices $d_\ell (\ell = 1, \dots, n)$, the integral $\int \Omega_K$ over the whole two-dimensional space reads $\int_{\mathbf{R}^2} \Omega_K = \pi Q = \pi \sum_{\ell=1}^n d_\ell Q_\ell$ where Q_ℓ are integers-valued topological charges. Hence, $\Omega_K(x)/\pi$ is identified with the density of the topological charge (up to the weight due to the index d_ℓ). Then, for the large Wilson loop compared with the typical size of the instanton, $\int_S \Omega_K(x)/\pi$ in the NAST counts the number of instantons minus anti-instantons which are contained inside the area $S \subset \mathbf{R}^2$ bounded by the loop C . Thus, the expectation value $\langle \exp(i \int_S \Omega_K) \rangle_{NLSM} = Z_{NLSM}^{-1} \int [d\mu(w, \bar{w})] \exp(-S_{NLSM}[w, \bar{w}]) \exp(i \int_S \Omega_K)$ is calculated by summing over all the possibilities of instanton and anti-instanton configurations or integration over the instanton moduli. For the quark in the fundamental representation ($d_1 = 1, d_2 = d_3 = \dots = d_n = 0$), this is easily performed as follows. Especially, in the $SU(3)$ case with $[1, 0]$, ξ is independent of w_3 , so, w_3 is redundant in the F_2 NLSM. Hence, it suffices to consider the CP^2 NLSM for the fundamental quark. If we put $w_2 = 0$, then $\Omega_K = i(1 + |w_1|^2)^{-2} dw_1 \wedge d\bar{w}_1$. Similarly, if $w_1 = 0$, then $\Omega_K = i(1 + |w_2|^2)^{-2} dw_2 \wedge d\bar{w}_2$. For a polynomial $w_\alpha = w_\alpha(z)$ in $z = x + iy$ with an order n , we find an instanton charge, $\int \Omega_K = \pi Q$, $Q = n$. This is the same situation as that encountered in $SU(2)$, $\int \Omega_K = 2j\pi Q$ ($j = 1/2$) [6, 7]. Thus the Wilson loop is estimated by the instanton calculus. In fact, the dilute instanton gas approximation leads to the area law for the Wilson loop, see [6]. This calculation is improved by including fluctuations from the instanton solutions and this issue will be discussed in detail elsewhere. In the paper

[9] we have applied the large N expansion to obtain the systematic result. At least in the leading order in $1/N$ expansion, we have obtained the result in consistent with instanton calculus for Wilson loop.

Finally, some comments are in order.

1) We want to emphasize that we have actually "derived" the area law of the Wilson loop by calculating the contributions from magnetic monopole. This is based on the combination of the non-Abelian Stokes theorem (NAST) and the reformulation of Yang-Mills theory within the framework of perturbative deformation. In other words, we have proposed a picture in which our framework is self-consistent.

Of course, we have obtained these results within the framework in which the quantum fluctuations around the magnetic monopole are treated in the perturbative way (which we called the perturbative deformation). However, this is the first analytical result of having derived the area law from the first principle by taking into account magnetic monopole in "3+1" dimensions. Our results should be regarded as the monopole dominance in the weak sense. That is to say, the magnetic monopole contribution is enough to derive the area law. The monopole dominance can be a sufficient condition for quark confinement, even if it is not a necessary condition

2) It should be remarked that the maximal stability group $\tilde{H}$ is in general different from the residual gauge group H for partial gauge fixing $G \rightarrow H$ and that $\tilde{H}$ is larger than H , i.e., $H \subset \tilde{H}$. For the coset, therefore, we have $G/\tilde{H} \subset G/H$. For $G = SU(3)$, $\tilde{H} = U(2)$ or $U(1) \times U(1)$ depending on the representation in question. By making use of the NAST, the Wilson loop is rewritten in terms of the $G/\tilde{H}$ variables. On the other hand, after partial gauge fixing $G \rightarrow H$, the gauge-fixing part is rewritten into the NLSM in terms of the G/H variables. So, the NLSM involves redundant variables, since $G/\tilde{H} \subset G/H$. Thus our claim is that the degrees of freedom which are directly responsible to the area law are given by $G/\tilde{H}$ rather than G/H . For the fundamental representation, $G/\tilde{H} = SU(3)/U(2) = CP^2$ and CP^{N-1} model has local gauge invariance $U(1)$ for any $N \geq 2$. In this sense, we can adopt even $H = U(2)$ in

the partial gauge fixing at least for the fundamental quark. In this case, $G/\tilde{H} = G/H$. Thus, the resultant theory after partial gauge fixing is not necessarily a $U(2)$ gauge theory and we don't intend to claim the equality of the string tensions between $U(2)$ and $SU(3)$. Rather, the two string tensions for $H = U(2)$ and $H = U(1) \times U(1)$ should coincide with each other.

3) Baryons are also important physical objects from the viewpoint of color confinement in the $SU(3)$ gauge theory as well as mesons. In the MA gauge, the remaining gauge symmetry is the local $U(1)^2$ symmetry and the global Weyl symmetry. The Weyl symmetry is a remanant of $SU(3)$, owing to which one will find only the color-singlet combination of quarks. In the presentation of our paper, however, there is no Weyl symmetry in a "manifest" form, because we can select the reference state from any of three states and we have selected a special highest weight vector denoted by Λ for the fundamental representation. Once a state is selected, one complex coordinate in F_2 becomes irrelevant in the expectation value of the Wilson loop and the relevant space becomes CP^2 which is spanned by other two complex coordinates in F_2 . The change of reference state causes only the change of irrelevant coordinate and the Wilson loop has the same expectation value. Therefore, the Weyl symmetry is easily recovered by averaging over all possible choices of the highest weight vector. The details will be given elsewhere.

Acknowledgments

This work is supported in part by the Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture (No.10640249).

References

- [1] Y. Nambu, Phys. Rev. D 10, 4262-4268 (1974).
G. 't Hooft, in: High Energy Physics, edited by A. Zichichi (Editorice Compositori, Bologna, 1975).
S. Mandelstam, Phys. Report 23, 245-249 (1976).

- [2] G. 't Hooft, Nucl. Phys. B 190 [FS3], 455-478 (1981).
A. Kronfeld, G. Schierholz and U.-J. Wiese, Nucl. Phys. B 293, 461-478 (1987).
A. Kronfeld, M. Laursen, G. Schierholz and U.-J. Wiese, Phys. Lett. B 198, 516-520 (1987).
- [3] T. Suzuki, hep-lat/9506016.
R.W. Haymaker, hep-lat/9510035.
A. DiGiacomo, hep-lat/9802008; hep-th/9603029.
M.I. Polikarpov, hep-lat/9609020. M.N. Chernodub and M.I. Polikarpov, hep-th/9710205.
G.S. Bali, hep-ph/9809351.
- [4] N. Seiberg and E. Witten, hep-th/9407087, Nucl. Phys. B 426, 19-52 (1994).
- [5] K.-I. Kondo, hep-th/9709109, Phys. Rev. D 57, 7467-7487 (1998).
- [6] K.-I. Kondo, hep-th/9801024, Phys. Rev. D 58, 105019 (1998).
- [7] K.-I. Kondo, hep-th/9805153, Phys. Rev. D 58, 105016 (1998).
- [8] K.-I. Kondo, hep-th/9904045.
- [9] K.-I. Kondo and Y. Taira, hep-th/9911242.
- [10] D.I. Diakonov and V.Yu. Petrov, Phys. Lett. B 224, 131-135 (1989).
D. Diakonov and V. Petrov, hep-th/9606104.
- [11] D.H. Feng, R. Gilmore and W.-M. Zhang, Rev. Mod. Phys. 62, 867-927 (1990).
- [12] A.M. Perelomov, Phys. Report 146, 135-213 (1987).
- [13] L. Faddeev and A.J. Niemi, hep-th/9807069; Phys. Rev. Lett. 82, 1624-1627 (1999).
L. Faddeev and A.J. Niemi, hep-th/9812090, Phys. Lett. B449, 214-218 (1999).
- [14] F. Wilczek and A. Zee, Phys. Rev. Lett. 52, 2111 (1984).
- [15] M. Quandt and H. Reinhardt, hep-th/9707185, Int. J. Mod. Phys. A13, 4049-4076 (1998).
- [16] K.-I. Kondo and T. Shinohara, in preparation.